

Benzofuran-, benzothiophene-, indazole- and benzisoxazole- quinones: excellent substrates for NAD(P)H:quinone oxidoreductase 1

**Jeffery J. Newsome,^a Mary Hassani,^b Elizabeth Swann,^a Jane M. Bibby,^a
Howard D. Beall^b and Christopher J. Moody^{a,c,*}**

^aDepartment of Chemistry, University of Exeter, Stocker Road, Exeter EX4 4QD, U.K.

^bDepartment of Biomedical and Pharmaceutical Sciences, The University of Montana, 32
Campus Drive #1552, Missoula, MT 59812-1552, U.S.A.

^cSchool of Chemistry, University of Nottingham, University Park, Nottingham NG7 2RD,
U.K.

Abstract

A series of heterocyclic quinones based on benzofuran, benzothiophene, indazole and benzisoxazole has been synthesized, and evaluated for their ability to function as substrates for recombinant human NAD(P)H:quinone oxidoreductase (NQO1), a two-electron reductase upregulated in tumor cells. Overall, the quinones are excellent substrates for NQO1, approaching the reduction rates observed for menadione.

1. Introduction

Quinones are widespread in nature,¹ and participate in important biological redox processes. For example, the ubiquinones act as electron-transfer agents in the respiratory chain, and the heterocyclic pyrroloquinolinequinone (coenzyme PQQ) is a redox co-factor. Indeed heterocyclic derivatives form an important subset of quinones,² often possessing potent

* Corresponding author: fax +44 115 951 3564; email c.j.moody@nottingham.ac.uk

biological activity, for example as phosphatase inhibitors.³ The best known heterocyclic quinone is the clinically used cancer therapeutic agent, the indolequinone natural product mitomycin C (MMC) **1**,⁴⁻⁷ although other heterocyclic quinones such as streptonigrin **2** have also been widely studied.⁸

Given the biological importance of quinone reduction, our own studies have focused on the two-electron reduction of quinones by the enzyme NAD(P)H:quinone oxidoreductase 1 (NQO1, DT-diaphorase, QR1).⁹⁻¹² In particular, we have investigated a broad range of indolequinone based substrates and inhibitors,¹³⁻¹⁷ the quinoline-5,8-dione system found in streptonigrin,¹⁸ and both benzimidazole-4,7-diones **3** and benzothiazole-4,7-diones **4** (Figure 1).¹⁹ Others have also studied benzimidazole quinones as analogues of MMC,²⁰ whilst 5-undecyl-6-hydroxybenzothiazole-4,7-dione (UHDBT), an analogue of ubiquinone, has been investigated as an inhibitor of electron transport by binding to cytochrome *bc*₁.²¹




Figure 1. Some heterocyclic quinones.


Despite the aforementioned studies, we sought to extend the range of heterocyclic quinones studied as substrates for NQO1, focusing on benzofurans, benzothiophenes, and indazoles. Furoquinones are quite well described, and although most of their naturally occurring compounds are naphthofuranquinones,¹ relatively simple derivatives such as acamelin **5** are known.²² Benzothiophene quinones are less common, although caldariellaquinone **6** appears to fulfil an important redox role in thermophilic and acidophilic archaea of the *Sulfolobus* genus that lack ubiquinones as electron-transfer agents.^{23,24} On the other hand, indazolequinones are somewhat rarer, although compounds such as **7** have been prepared as MMC analogues,²⁵ whilst others have been investigated as substrates for carbonyl reductase.²⁶ Therefore in an attempt to widen the group of NQO1 substrates/inhibitors, and to probe further the active site of the enzyme, we have explored a new series of heterocyclic quinones based on benzofuran, benzothiophene, indazole and benzisoxazole.

2. Results and Discussion

2.1 Chemistry


In order to make meaningful comparisons with the more widely studied indolequinones, we initially elected to investigate relatively simple 5-methoxy-heterocyclic quinones in the benzofuran and benzothiophene series. The synthesis of quinone **11** started with the known benzofuran **8**, readily prepared from benzoquinone in a Nenitzescu type reaction.²⁷ Methylation gave the known 5-methoxy derivative **9**, nitration of which gave a mixture of the desired 4-nitro compound **10** along with its 6-nitro isomer in excellent overall yield, but in a 1:2 ratio. Although nitrobenzofuran **10** could be isolated, it was more convenient to reduce the mixture of nitro compounds to the corresponding amines, reduce the ketone with sodium

borohydride and then oxidize with Fremy's salt and purify the desired quinone **11** at the final stage (Scheme 1).




Scheme 1. Reagents and conditions: a, KH, MeI, DMF, rt (42%); b, fuming HNO₃, AcOH, rt (81% as 1:2 mixture of 4- and 6-nitro isomers); c, Sn, HCl, EtOH, reflux; d, NaBH₄, MeOH, rt; e, Fremy's salt, NaH₂PO₄, aq acetone, rt (20% over 3 steps).

The isomeric benzofuranquinone **13**, with the alcohol group at the 2-position, was synthesized from the known benzofuran **12**, readily available from 4-methoxyphenol by carbene O-H insertion, and cyclization.²⁸ The sequence of nitration, reduction and oxidation to the quinone was carried out without purification of the intermediate compounds and delivered the pure benzofuranquinone **13** in 53% over the four steps (Scheme 2).




Scheme 2. Reagents and conditions: a, HNO₃, AcOH, rt; b, Sn, HCl, EtOH, rt; c, LiAlH₄, THF, 0 °C; d, Fremy's salt, NaH₂PO₄, aq acetone, rt (53% over 4 steps).

In the benzothiophene series, we started with the known 5-methoxy-2-methylbenzothiophene **14** (Scheme 3).²⁹ Formylation under Vilsmeier conditions gave the desired 3-aldehyde **15**, but in poor yield (20%), the major product being the unwanted 4-formyl compound. Nevertheless, subjecting aldehyde **15** to the usual sequence of nitration, reduction, and final oxidation to the quinone, provided the benzothiophenequinone **18** in reasonable overall yield (Scheme 3).




Scheme 3. *Reagents and conditions:* a, POCl₃, DMF, CH₂Cl₂, 0 to 50 °C (20%, plus 40% of 4-formyl isomer); b, HNO₃, AcOH, rt (61%); c, Sn, HCl, EtOH, reflux (71%); d, NaBH₄, MeOH, rt; e, Fremy's salt, NaH₂PO₄, aq acetone, rt (41% over 2 steps).

By analogy with the conversion of benzofuran **12** into benzofuranquinone **13**, the known starting material,³⁰ ethyl 5-methoxy-3-methylbenzothiophene-2-carboxylate **19**, was converted into benzothiophenequinone **23** by the now familiar sequence of nitration, reduction and final oxidation to the quinone (Scheme 4).



Scheme 4. Reagents and conditions: a, HNO₃, AcOH, rt (74%); b, Sn, HCl, EtOH, reflux (86%); d, LiAlH₄, THF, rt (80%); d, Fremy's salt, NaH₂PO₄, aq acetone, rt (67%).


A range of 3-unsubstituted benzothiophenequinones was also prepared as shown in Scheme 5. Reaction of 3-methoxycinnamic acid with thionyl chloride and methanol readily provided the known benzothiophene-2-carboxylate starting material **24**.³¹ Nitration, followed by hydrogenation over Pd/C resulted in reduction of the nitro group with concomitant hydrogenolysis of the chlorine, and gave the 4-aminobenzothiophene **26**, which was either directly oxidized to quinone **27**, or converted into quinone **29** by initial reduction of the ester (Scheme 5). Benzothiophenequinones **31-33** were also prepared by standard transformations as shown in Scheme 5.



Scheme 5. Reagents and conditions: a, HNO₃, AcOH, reflux (87%); b, H₂, Pd/C, MeOH-THF (83%); c, Fremy's salt, NaH₂PO₄, aq acetone, rt (75%); d, LiAlH₄, THF, rt (78%); e, Fremy's salt, NaH₂PO₄, aq acetone, rt (88%); f, MnO₂, CH₂Cl₂, reflux (38%); g, Fremy's salt, NaH₂PO₄, aq acetone, rt (79%); h, Ac₂O, pyridine, rt (69%); i, MsCl, Et₃N, CH₂Cl₂, rt; j, LiAlH₄, THF, rt (5% over 2 steps);


The synthesis of indazolequinones started from the dimethoxyindazole **35**, itself prepared by the literature method from 3,5-dimethoxyacetophenone **34** by electrophilic amination with bis(trichloroethyl) azodicarboxylate followed by zinc reduction and cyclization.³² Nitration, methylation, reduction and oxidation delivered the required indazolequinone **39** (Scheme 6).

The methyl group was further functionalized by radical bromination and conversion into the 3-hydroxymethylindazolequinone **40** (Scheme 6).



Scheme 6. Reagents and conditions: a, Cl₃CCH₂O₂CN=NCO₂CH₂Cl₃, BF₃•OEt₂, CH₂Cl₂, rt; b, Zn, AcOH, rt (35% over 2 steps); c, HNO₃, AcOH, rt (81%); d, MeI, KOH, DMSO, rt (49%); e, Sn, HCl, EtOH, reflux (96%); f, Fremy's salt, NaH₂PO₄, aq acetone, rt (78%); g, NBS, AIBN, CCl₄; h, AgNO₃, aq acetone (36% over 2 steps).

Finally an example of a benzisoxazolequinone was prepared. 2-Hydroxy-5-methoxyacetophenone **41** was converted into its oxime **42** that underwent facile intramolecular Mitsunobu reaction to provide the benzisoxazole **43** in quantitative yield. Thereafter, nitration, reduction and oxidation as before gave the required benzisoxazolequinone **45** (Scheme 7)



Scheme 7. Reagents and conditions: a, $\text{NH}_2\text{OH}\cdot\text{HCl}$, EtOH, pyridine, rt (98%); b, DIAD, Ph_3P , THF, rt (100%); c, HNO_3 , AcOH, rt (91%); d, Sn, HCl, EtOH, reflux; e, Fremy's salt, NaH_2PO_4 , aq acetone, rt (20% over 2 steps).

Reduction potentials were measured for benzofuran-, benzothiophene- and indazole-quinones **13**, **18**, **23**, **39** and **40** using cyclic voltammetry in DMF as solvent with tetra-*n*-butylammonium tetrafluoroborate as supporting electrolyte as previously described.¹³ The E_{redox} values, with reference to ferrocene (Fc), are shown in Figure 2; values for the related indolequinones **46** and **47** are also shown. The data show that whilst the indazole quinone **39** has a similar redox potential to the indolequinones (E_{redox} v. Fc -1.20 to -1.40V), the other heterocyclic quinones are considerably easier to reduce.





Figure 2. E_{redox} values (v. Fc) for benzimidazole- and benzothiazole-quinones **11** and **25** compared to related indolequinones **46** and **47**.^{13,14}

2.2 Enzyme Studies

The new heterocyclic quinones were evaluated for their ability to act as substrates for NQO1. We used two assays for studying quinone metabolism by recombinant human NQO1 based on HPLC and spectrophotometry. The former HPLC system is capable of quantifying both NADH oxidation and quinone reduction, and gives average rates of reduction over a 30 – 40 minute period.^{33,13} Quinone reduction is reversible due to redox cycling of the hydroquinone, so results (Table 1) are reported as $\mu\text{mol NADH oxidized min}^{-1} \text{mg}^{-1}$ NQO1. The alternative spectrophotometric method uses cytochrome *c* as the terminal electron acceptor and gives initial rates of reduction that are generally higher than the HPLC method.¹⁸ Nevertheless the relative order of metabolism is essentially the same with the two methods, and with the

exception of Entries 1, 2, 5 and 6, both methods were used to enable reliable comparison between the new heterocyclic quinones.

Table 1. Metabolism of heterocyclic quinones by recombinant human NQO1.



Entry	Ring	Cpd	X	R ²	R ³	NQO1 (ave) ^a μmol/min/mg	NQO1 (init) ^a μmol/min/mg
1	A	46	NMe	Me	CH ₂ OH	1.25 ± 0.03 ^b	<i>nd</i> ^c
2	A	47	NMe	CH ₂ OH	Me	2.49 ± 1.27 ^d	<i>nd</i> ^c
3	A	11	O	Me	CH(OH)Me	87.9±25.6	585±32
4	A	13	O	CH ₂ OH	Me	80.8±6.7	1025±54
5	A	18	S	Me	CH ₂ OH	<i>nd</i> ^c	236±35
6	A	23	S	CH ₂ OH	Me	<i>nd</i> ^c	468±68
7	A	27	S	CO ₂ Me	H	53.9±3.8	555±71
8	A	29	S	CH ₂ OH	H	63.8±11.7	927±89
9	A	31	S	CHO	H	27.4±5.7	897±64
10	A	32	S	CH ₂ OAc	H	24.5±7.9	900±86
11	A	33	S	Me	H	5.51±0.87	553±66
12	B	39	NMe	-	Me	19.6±5.3	55.3±5.1
13	B	40	NMe	-	CH ₂ OH	38.1±7.1	247±14
14	B	45	O	-	Me	27.6±3.2	290±28

^a Ave and init refer to average and initial rates of metabolism as measured by the HPLC and spectrophotometric assays, respectively; average rates are determined from irreversible NADH oxidation whereas initial rates are determined from cytochrome *c* reduction. For reference, initial rate for menadione reduction was 1225 ± 15 μmol/min/mg. ^b Ref¹³; ^c *nd* = not determined; ^d Ref¹⁴.

The new quinones are all excellent substrates for rhNQO1. In the benzofuran and benzothiophene series, reduction rates were higher when the hydroxyalkyl substituent was at the C-2 position rather than C-3 (Table 1). As with the benzimidazole- and benzothiazole-quinones,¹⁹ all of the new quinones were much better substrates for NQO1 than the widely studied indolequinones,^{13,33,14} as seen by comparison with indolequinones **46** and **47** included in Table 1 for comparison. In fact, the reduction rates for benzofuran- and benzothiophene-quinones **13**, **29**, **31** and **32** approach the initial reduction rate observed for menadione ($1225 \pm 15 \mu\text{mol}/\text{min}/\text{mg}$),¹⁹ a simple naphthoquinone that has been used to measure activity of the enzyme, making these compounds some of the best NQO1 substrates seen to date.

The results presented here complement our previous work on bioreductive activation of indolequinone antitumor agents by NQO1. Novel heterocyclic quinones have been synthesized, characterized and studied biologically as substrates for recombinant human NQO1. These data add to our understanding of the structural requirements for efficient metabolism by the quinone reductase enzyme.

3. Experimental Section

3.1 General Chemistry Experimental Details


Commercially available reagents were used throughout without purification unless otherwise stated. Light petroleum refers to the fraction with bp 40 – 60 °C and was distilled before use. Ether refers to diethyl ether. Reactions were routinely carried out under a nitrogen or argon atmosphere. Analytical thin layer chromatography was carried out on aluminum-backed plates coated with Merck Kieselgel 60 GF₂₅₄, and visualized under UV light at 254 and/or 360

nm. Flash chromatography was carried out using Merck Kieselgel 60 H silica or Matrex silica 60. Fully characterized compounds were chromatographically homogeneous.

Infrared spectra were recorded in the range 4000 – 600 cm^{-1} using FT spectrometers. NMR spectra were carried out at 300 and 400 MHz (^1H frequencies, corresponding ^{13}C frequencies are 75 and 100 MHz). Chemical shifts are quoted in ppm with TMS as internal standard. J values are recorded in Hz. In the ^{13}C spectra, signals corresponding to CH, CH_2 or CH_3 groups, as assigned from DEPT, are noted; all others are C. High and low resolution mass spectra were recorded on a Micromass GCT TDF High Resolution mass spectrometer, or at the EPSRC Mass Spectrometry Service (Swansea).


3.2 Synthesis of Benzofuranquinones

3-Acetyl-5-methoxy-2-methylbenzofuran **9**




3-Acetyl-5-hydroxy-2-methylbenzofuran **8**²⁷ (1.62 g, 8.5 mmol) in DMF (30 mL) was added to a stirring suspension of potassium hydride (0.61 g, 15.2 mmol) in DMF (50 mL) at 0 °C. The mixture was stirred at room temperature for 45 min. Iodomethane (1.80 g, 12.7 mmol) was added dropwise at 0 °C and the mixture allowed to warm to room temperature. After 2 h saturated ammonium chloride solution was added and the mixture extracted with ethyl acetate. The ethyl acetate layer was washed thoroughly with hydrochloric acid (1 M), dried (MgSO_4) and concentrated. The crude product was purified by chromatography eluting with dichloromethane to yield the *title compound* as a colorless solid (0.74 g, 42%), mp 69-70 °C (lit.,³⁴ mp 72 °C); δ_{H} (300 MHz; CDCl_3) 7.48 (1 H, d, J 2.6, 4-H), 7.33 (1 H, d, J 8.9, 7-H), 6.89 (1 H, dd, J 8.9, J 2.6, 6-H), 3.88 (3 H, s, OMe), 2.76 (3 H, s, Me), 2.62 (3 H, s, Me).

3-Acetyl-5-methoxy-2-methyl-4-nitrobenzofuran **10**



To a solution of 3-acetyl-5-methoxy-2-methylbenzofuran **9** (0.217 g, 1.1 mmol) in acetic acid (4 mL) cooled to 0 °C was added a mixture of fuming nitric acid (0.5 mL) and acetic acid (2 mL). The mixture was stirred at room temperature for 2 h and then poured on to an ice water mixture and the precipitate obtained was filtered off and dried. NMR analysis of the mixture showed a 2:1 ratio of 6-nitro and 4-nitro products (0.21 g, 81%), used without further purification. Small quantities of each isomer were obtained by preparative TLC (dichloromethane elution) and were identified by ¹H NMR spectroscopy: *3-acetyl-5-methoxy-2-methyl-4-nitrobenzofuran 10*, δ_{H} (300 MHz; CDCl₃) 7.50 (1 H, d, *J* 9.0, ArH), 7.03 (1 H, d, *J* 9.0 ArH), 3.93 (3 H, s, OMe), 2.74 (3 H, s, Me), 2.49 (3 H, s, Me); *3-acetyl-5-methoxy-2-methyl-6-nitrobenzofuran*, δ_{H} (300 MHz; CDCl₃) 7.97 (1 H, s, ArH), 7.76 (1 H, s, ArH), 4.02 (3 H, s, OMe), 2.83 (3 H, s, Me), 2.62 (3 H, s, Me).


3-(1-Hydroxyethyl)-5-methoxy-2-methylbenzofuran-4,7-dione **11**



To a mixture of 3-acetyl-5-methoxy-2-methyl-4-nitrobenzofuran **10** and its 6-nitro isomer (1:2 ratio) (0.80 g, 3.4 mmol) in ethanol (100 mL) were added tin powder (4.00 g, 34.0 mmol) and hydrochloric acid (3 M; 40 mL). The mixture was heated under reflux for 30 min. Upon cooling the solution was decanted from the tin and neutralized with saturated aqueous sodium hydrogen carbonate. The mixture was extracted with ethyl acetate and the organic layer dried (MgSO₄) and concentrated, to yield a mixture of 4- and 6-amino derivatives (0.56 g) that was

used directly. Sodium borohydride (0.10 g, 2.6 mmol) was added to a stirred solution of the above mixture (0.56 g, 2.6 mmol) in methanol (50 mL). After 20 min, acetone (10 mL) was added. The solvent was removed and the residue dissolved in ethyl acetate and washed with water. The organic layer was dried (MgSO₄) and concentrated. To a solution of the crude product in acetone (40 mL) was added a solution of potassium nitrosodisulfonate (1.00 g, 3.7 mmol) in sodium dihydrogen phosphate buffer (0.3 M; 40 mL). The mixture was stirred at room temperature for 1 h. The acetone was removed *in vacuo*, and the resulting residue was extracted with dichloromethane and washed with water. The organic layer was dried (Na₂SO₄) and concentrated. The crude material was purified by chromatography eluting with dichloromethane/ethyl acetate (19:1) and recrystallized (dichloromethane/ether) to yield the *title compound* (0.052 g, 20%, over 3 steps based on the 4-nitro starting material), as an orange crystalline solid, mp 157-158 °C; λ_{\max} (MeOH)/nm 436 (log ϵ 3.18), 320 (3.66), 260 (3.98); ν_{\max} (KBr)/cm⁻¹ 3414, 3060, 2990, 2915, 1683, 1657, 1607, 1587; δ_{H} (300 MHz; CDCl₃) 5.76 (1 H, s, 6-H), 4.81 (1 H, m, CHMe), 3.85 (3 H, s, OMe), 3.78 (1 H, d, *J* 10.7 Hz, OH), 2.38 (3 H, s, Me), 1.45 (3 H, d, *J* 6.7 Hz, CHMe); δ_{C} (100 MHz; CDCl₃) 179.2 (C), 175.4 (C), 159.8 (C), 153.4 (C), 150.6 (C), 125.0 (C), 124.8 (C), 105.7 (CH), 62.4 (CH), 56.9 (Me), 24.4 (Me), 12.2 (Me).

2-Hydroxymethyl-5-methoxy-3-methylbenzofuran-4,7-dione 13




To a solution of methyl 5-methoxy-3-methylbenzofuran-2-carboxylate **12**²⁸ (0.20 g, 0.91 mmol) in acetic acid (6 mL), cooled to -10°C was added a mixture of concentrated nitric acid (1 mL) and acetic acid (4 mL). The mixture was stirred at room temperature for 1 h. The reaction mixture was poured in an ice/water mixture and the resulting precipitate filtered off

and dried. The crude material was obtained as a 2.8:1 ratio of 4- and 6-nitro products and used directly in the next step. To a suspension of the above mixture in ethanol (30 mL) were added tin powder (0.48 g, 4.0 mmol) and hydrochloric acid (3 M; 7 mL). The mixture was stirred at room temperature for 2 h. The solution was decanted from the excess tin and neutralized with saturated aqueous sodium hydrogen carbonate. The suspension obtained was added to an equal volume of water. The mixture was extracted with ethyl acetate. The organic layer was dried (MgSO_4) and concentrated. The crude material was used directly in the next step without purification.

To a suspension of lithium aluminum hydride (0.091 g, 2.4 mmol) in THF (10 mL) at 0°C was added a solution of the above mixture of benzofurans in THF (10 mL) and the reaction was stirred for 15 min. The reaction mixture was quenched by the addition of water (0.2 mL), sodium hydroxide (1 M; 0.2 mL) and silica gel (2 g). The granular precipitate was filtered off through a pad of Celite. The filtrate was dried (MgSO_4) and concentrated *in vacuo* to give the alcohol that was used directly in the next step without purification, or characterization. To a solution of the benzofuran-2-methanol in acetone (50 mL) was added a solution of potassium nitrosodisulfonate (0.64 g, 2.4 mmol) in sodium dihydrogen phosphate buffer (0.3 M; 50 mL). The mixture was stirred at room temperature for 1 h. The excess acetone was removed *in vacuo*. The resulting residue was extracted with dichloromethane and washed with water. The organic layer was dried (Na_2SO_4) and concentrated. The crude product was purified by chromatography eluting with ethyl acetate/dichloromethane (1:4) and recrystallized (ethyl acetate) to yield the *title compound* as an orange solid (0.108 g, 53%, over 4 steps); mp 205-206 $^\circ\text{C}$; (Found: C, 58.4; H, 4.5. $\text{C}_{11}\text{H}_{10}\text{O}_5 + 0.2\text{H}_2\text{O}$ requires C, 58.5; H, 4.6%); (Found: M^+ , 222.0528. $\text{C}_{11}\text{H}_{10}\text{O}_5$ requires 222.0528); ν_{max} (KBr)/ cm^{-1} 3452, 3057, 2939, 1691, 1660, 1610, 1587; δ_{H} (300 MHz; CDCl_3) 5.80 (1 H, s, 6-H), 4.69 (2 H, d, J 4.0 Hz, CH_2OH), 3.86 (3 H, s, OMe), 2.31 (3 H, s, Me), 1.98 (1 H, br s, OH); δ_{C} (100 MHz; acetone) 177.8 (C), 175.3 (C),

160.3 (C), 156.6 (C), 149.2 (C), 125.4 (C), 116.7 (C), 105.7 (CH), 56.4 (Me), 54.0 (CH₂), 7.6 (Me); *m/z* (EI) 222 (M⁺, 35%), 193 (13), 151 (23), 109 (25), 69 (100).

The intermediate nitro compound methyl 5-methoxy-3-methyl-4-nitrobenzofuran-2-carboxylate, can be isolated and characterized.




To a solution of methyl 5-methoxy-3-methylbenzofuran-2-carboxylate **12** (0.130 g, 0.6 mmol) in acetic acid (5 mL), cooled to -10°C was added a mixture of concentrated nitric acid (0.5 mL) and acetic acid (3 mL). The mixture was stirred at room temperature for 1 h. The reaction mixture was poured in an ice/water mixture and the resulting precipitate filtered off and dried.

The crude material was obtained as a 3:1 ratio of 4- and 6-nitro products (0.12 g, 80%).

Purification by chromatography eluting with dichloromethane gave *methyl 5-methoxy-3-methyl-4-nitrobenzofuran-2-carboxylate*, mp 146-149 °C; ν_{\max} (KBr)/cm⁻¹ 3103, 2996, 2945, 1706, 1629, 1588; δ_{H} (300 MHz; CDCl₃) 7.64 (1 H, d, *J* 9.1 Hz, ArH), 7.20 (1 H, d, *J* 9.1 Hz, ArH), 3.99 (3 H, s, OMe), 3.96 (3 H, s, OMe), 2.47 (3 H, s, Me); δ_{C} (75 MHz; CDCl₃) 160.0 (C), 148.8 (C), 147.6 (C), 143.3 (C), 133.6 (C), 123.0 (C), 121.5 (C), 115.1 (CH), 113.7 (CH), 57.7 (Me), 52.4 (Me), 8.9 (Me).

3.3 Synthesis of Benzothiophenequinones


5-Methoxy-2-methylbenzothiophene-3-carboxaldehyde **15**



To a mixture of dry DMF (19.6 mL, 169.40 mmol) and dry dichloromethane (9.1 mL) was added dropwise phosphorus oxychloride (16.4 mL, 169.40 mmol) at 0 °C. The ice bath was

removed. The reaction mixture was stirred for an additional 30 min and cooled in an ice bath. 5-Methoxy-2-methylbenzothiophene **14**²⁹ (3.02 g, 16.94 mmol) was added portionwise to the solution over 5 min at 0 °C. The reaction mixture was stirred and heated to 50 °C overnight, poured into an ice-cold aqueous sodium hydroxide (1 M). The mixture was extracted with dichloromethane, dried over MgSO₄, filtered, and concentrated. The residue obtained was purified by chromatography, eluting with dichloromethane, to yield the *title compound* (390 mg, 20%) as a yellow oil; (Found: MH⁺, 207.0477. C₁₁H₁₀O₂S + H requires 207.0480); ν_{\max} (film)/cm⁻¹ 2957, 2930, 2834, 1735, 1670, 1597, 1559, 1455, 1413, 1351, 1270, 1247, 1224, 1151; δ_{H} (300 MHz; CDCl₃) 10.31 (1 H, s, CHO), 8.14 (1 H, d, *J* 2.5, 4-H), 7.60 (1 H, d, *J* 9.0, 7-H), 7.00 (1 H, dd, *J* 9.0, 2.5, 6-H), 3.88 (3 H, s, OMe), 2.89 (3 H, s, Me); δ_{C} (75 MHz; CDCl₃) 184.0 (CH), 159.6 (C), 159.3 (C), 138.9 (C), 130.5 (C), 129.3 (C), 122.6 (CH), 116.0 (CH), 106.4 (CH), 56.1 (Me), 14.8 (Me); *m/z* (CI) 207 (MH⁺, 100%), 179 (10); and the *4-formyl derivative* (809 mg; 40 %) as a beige solid; mp 119-120 °C; (Found: MH⁺, 207.0477. C₁₁H₁₀SO₂ + H requires 207.0480); ν_{\max} (KBr)/cm⁻¹ 3421, 2910, 2846, 1661, 1566, 1454, 1430, 1314, 1243, 1183, 1131, 1071; δ_{H} (300 MHz; CDCl₃) 10.68 (1 H, s, CHO), 8.09 (1 H, br s, 3-H), 7.86 (1 H, d, *J* 8.8, ArH), 6.96 (1 H, d, *J* 8.8, ArH), 3.97 (3 H, s, OMe), 2.63 (3 H, s, Me); δ_{C} (100 MHz; CDCl₃) 190.7 (CH), 162.3 (C), 148.0 (C), 139.9 (C), 134.1 (C), 129.7 (CH), 122.7 (CH), 118.1 (C), 108.6 (CH), 55.8 (Me), 17.0 (Me); *m/z* (CI) 207 (MH⁺, 100%), 179 (10).


5-Methoxy-2-methyl-4-nitrobenzothiophene-3-carboxaldehyde **16**



To a solution of 5-methoxy-2-methylbenzothiophene-3-carboxaldehyde **15** (388 mg, 1.88 mmol) in acetic acid (3.2 mL), cooled to 0-5 °C was added a mixture of nitric acid (638 μ l,

9.42 mmol) in acetic acid (4.7 mL). The mixture was stirred at room temperature overnight. The reaction mixture was poured into ice/water, neutralized with a saturated aqueous sodium hydrogen carbonate, extracted with dichloromethane, dried over MgSO₄, filtered, and the filtrate evaporated under reduced pressure and azeotroped with toluene to remove the remaining acetic acid. The crude product obtained was purified by chromatography, eluting with dichloromethane, to yield the *title compound* (286 mg; 61 %) as a yellow solid; mp 205-207 °C; (Found: MH⁺, 252.0333. C₁₁H₉NO₄S + H requires 252.0331); ν_{\max} (KBr)/cm⁻¹ 2975, 2941, 2845, 2771, 1679, 1598, 1536, 1459, 1425, 1375, 1271, 1117, 1082; δ_{H} (300 MHz; CDCl₃) 10.05 (1 H, s, CHO), 7.80 (1 H, d, *J* 8.9, ArH), 7.66 (1 H, d, *J* 8.9, ArH), 3.97 (3 H, s, OMe), 2.90 (3 H, s, Me); δ_{C} (75 MHz; CDCl₃) 183.7 (CHO), 160.4 (C), 150.4 (C), 131.0 (C), 129.4 (C), 128.8 (C), 125.1 (CH), 111.4 (CH), 57.7 (Me), 16.4 (Me), one C unobserved; *m/z* (CI) 252 (MH⁺, 100%), 222 (65), 206 (30), 194 (10).


4-Amino-5-methoxy-2-methylbenzothiophene-carboxaldehyde **17**



To a suspension of 5-methoxy-2-methyl-4-nitrobenzothiophene-carboxaldehyde **16** (280 mg, 1.12 mmol) in ethanol (23 mL) was added tin powder (601 mg, 5.02 mmol) and hydrochloric acid (3 M; 8.1 mL). The mixture was stirred and heated under reflux for 1 h. Upon cooling, the reaction mixture was decanted from the excess of tin and neutralized with a saturated aqueous solution of NaHCO₃. The suspension obtained was filtered through Celite and extracted with ethyl acetate. The combined organic layer was dried over MgSO₄ and evaporated under reduced pressure to yield the *title compound* (175 mg, 71%) as a yellow solid; mp 88-90 °C; (Found: MH⁺, 222.0587. C₁₁H₁₁NO₂S + H requires 222.0589); ν_{\max} (KBr)/cm⁻¹ 3437, 2960, 2921, 2852, 1655, 1602, 1559, 1459, 1421, 1340, 1224, 1201; δ_{H} (300

MHz; CDCl₃) 10.08 (1 H, s, CHO), 6.96 (1 H, d, *J* 8.5, ArH), 6.91 (1 H, d, *J* 8.5, ArH), 6.02 (2 H, br s, NH₂), 3.89 (3 H, s, OMe), 2.85 (3 H, s, Me); δ_C (100 MHz; CDCl₃) 184.7 (CH), 160.8 (C), 144.2 (C), 134.3 (C), 132.1 (C), 130.9 (C), 123.1 (C), 110.5 (CH), 108.6 (CH), 56.5 (Me), 14.9 (Me); *m/z* (CI) 222 (MH⁺, 100%), 206 (5).

3-Hydroxymethyl-5-methoxy-2-methylbenzothiophene-4,7-dione 18




Sodium borohydride (44 mg, 2.31 mmol) was added in one portion to a solution of 4-amino-5-methoxy-2-methylbenzothiophene-3-carboxaldehyde **17** (170 mg, 0.77 mmol) in dry methanol (21 mL) cooled at 0 °C. The reaction mixture was stirred at room temperature for 1 h. The reaction mixture was evaporated off and the crude product obtained was purified by chromatography, eluting with ethyl acetate-dichloromethane 1:1, to yield the unstable *amino alcohol intermediate* as a yellow solid that was used in the next step with no further purification.

To a solution of the *amino alcohol intermediate* dissolved in acetone (50 mL) was added a solution of potassium nitrosodisulfonate (907 mg, 3.32 mmol) in sodium dihydrogen phosphate buffer (0.3 M; 40 mL). The reaction was stirred at room temperature for 1 h. The excess acetone was removed *in vacuo*. The resulting residue was extracted with dichloromethane and the combined organic layer evaporated off. The residue obtained was purified by chromatography, eluting with dichloromethane, to yield the *title compound* (75 mg, 41%) as a yellow crystalline solid; mp 196-198 °C; (Found: MH⁺, 239.0372. C₁₁H₁₀O₄S + H requires 239.0378); λ_{max} (acetonitrile)/nm 200 (log ε 4.14), 232 (4.12), 288 (4.18), 348 (3.47), 416 (3.21); ν_{max} (KBr)/cm⁻¹ 3425, 2925, 2852, 1663, 1636, 1601, 1447, 1344, 1324, 1255, 1220, 1120, 1086, 1070; δ_H (300 MHz; CDCl₃) 5.96 (1 H, s, 6-H), 4.68 (2 H, s,


CH_2OH), 3.88 (3 H, s, OMe), 2.53 (3 H, s, Me); δ_{C} (100 MHz; CDCl_3) 179.5 (C), 178.1 (C), 160.1 (C), 144.4 (C), 142.9 (C), 138.4 (C), 137.0 (C), 108.1 (CH), 56.9 (Me), 56.5 (CH_2), 13.7 (Me); m/z (CI) 239 (MH^+ , 85%), 223 (50), 221 (100).

Ethyl 5-methoxy-3-methyl-4-nitrobenzothiophene-2-carboxylate **20**




To a solution of ethyl 5-methoxy-3-methylbenzothiophene-2-carboxylate **19**³⁰ (480 mg, 1.92 mmol) in acetic acid (3.2 mL), cooled to 0-5 °C was added a mixture of nitric acid (1.3 mL, 19.20 mmol) in acetic acid (4.8 mL). The mixture was stirred at room temperature overnight. The reaction mixture was dropped in ice/water, extracted with dichloromethane, dried over MgSO_4 , filtered, evaporated and azeotroped with toluene to remove the acetic acid. The crude product obtained was purified by chromatography, eluting with dichloromethane, to yield the *title compound* (417 mg, 74%) as a light yellow solid; mp 158-160 °C; (Found: MH^+ , 296.0589. $\text{C}_{13}\text{H}_{13}\text{NO}_5\text{S} + \text{H}$ requires 296.0592); ν_{max} (KBr)/ cm^{-1} 3422, 2988, 2939, 2850, 1711, 1608, 1527, 1446, 1370, 1297, 1247, 1193, 1174, 1151, 1051; δ_{H} (300 MHz; CDCl_3) 7.85 (1 H, d, J 8.9, ArH), 7.26 (1 H, d, J 8.9, ArH), 4.40 (2 H, q, J 7.1, OCH_2Me), 3.98 (3 H, s, OMe), 2.64 (3 H, s, Me), 1.41 (3 H, t, J 7.1, CH_2Me); δ_{C} (100 MHz; CDCl_3) 162.6 (C), 148.6 (C), 137.1 (C), 136.7 (C), 133.8 (C), 131.5 (C), 130.6 (C), 125.1 (CH), 113.3 (CH), 61.7 (CH_2), 57.3 (Me), 14.2 (Me), 12.1 (Me); m/z (CI) 296 (MH^+ , 100%), 279 (8), 266 (15).

Ethyl 4-amino-5-methoxy-3-methylbenzothiophene-2-carboxylate **21**



To a suspension of ethyl 5-methoxy-3-methyl-4-nitrobenzothiophene-2-carboxylate **20** (370 mg, 1.25 mmol) in ethanol (27 mL) was added tin powder (676 mg, 5.64 mmol) and hydrochloric acid (3 M; 9 mL). The mixture was stirred and heated under reflux for 1 h. Upon cooling, the reaction mixture was decanted from the excess of tin and neutralized with a saturated aqueous solution of NaHCO₃. The suspension obtained was filtered through Celite and extracted with ethyl acetate. The combined organic layer was dried over MgSO₄, filtered and evaporated under reduced pressure to yield the *title compound* (285 mg, 86%) as a yellow solid; mp 120-123 °C; (Found: MH⁺, 266.0844. C₁₃H₁₅NO₃S + H requires 266.0851); ν_{\max} (KBr)/cm⁻¹ 3460, 3368, 2968, 2925, 1705, 1621, 1536, 1471, 1440, 1332, 1259, 1209, 1178, 1144, 1094, 1055, 1009; δ_{H} (300 MHz; CDCl₃) 7.11 (1 H, d, *J* 8.6, ArH), 7.03 (1 H, d, *J* 8.6, ArH), 4.35 (2 H, q, *J* 7.0, OCH₂Me), 3.89 (3 H, s, OMe), 3.06 (3 H, s, Me), 1.40 (3 H, t, *J* 7.0, CH₂Me); NH₂ not observed; δ_{C} (100 MHz; CDCl₃) 165.7 (C), 145.5 (C), 143.3 (C), 136.7 (C), 136.5 (C), 130.5 (C), 127.3 (C), 115.3 (CH), 113.4 (CH), 63.1 (CH₂), 59.0 (Me), 18.1 (Me), 16.4 (Me); *m/z* (CI) 266 (MH⁺, 100%), 265 (40), 220 (10).


4-Amino-5-methoxy-3-methylbenzothiophene-2-methanol **22**



To a suspension of lithium aluminum hydride (162 mg, 4.23 mmol) in dry THF (6 mL) at 0 °C was added a solution of ethyl 4-amino-5-methoxy-3-methylbenzothiophene-2-carboxylate **21** (280 mg, 1.06 mmol) in dry THF (3 mL). The mixture was allowed to warm up to room temperature and stirred for 2 h. The mixture was cooled to 0 °C and quenched by the addition of water (0.5 mL), aqueous sodium hydroxide (1 M; 0.5 mL) and silica gel. The granular precipitate was filtered off through a pad of Celite. The filtrate was dried over MgSO₄, filtered and concentrated *in vacuo* to yield the *title compound* (188 mg, 80%) as an orange

solid; mp 114-115 °C; (Found: MH^+ , 224.0753. $\text{C}_{11}\text{H}_{11}\text{NO}_2\text{S} + \text{H}$ requires 224.0745); ν_{max} (KBr)/ cm^{-1} 3436, 3352, 3273, 2923, 2851, 1599, 1462, 1265, 1203, 1157, 1136, 1044, 1019; δ_{H} (300 MHz; CDCl_3) 7.12 (1 H, d, J 8.6, ArH), 6.90 (1 H, d, J 8.6, ArH), 4.75 (2 H, s, CH_2OH), 3.88 (3 H, s, OMe), 2.54 (3 H, s, Me); OH, NH_2 not observed; δ_{C} (100 MHz; CDCl_3) 142.3 (C), 134.8 (C), 131.7 (C), 131.1 (C), 127.5 (C), 126.8 (C), 110.6 (CH), 109.1 (CH), 56.9 (CH_2), 55.5 (Me), 13.4 (Me); m/z (CI) 224 (MH^+ , 60%), 223 (M^+ , 80), 206 (100), 194 (15).


2-Hydroxymethyl-5-methoxy-3-methylbenzothiophene-4,7-dione **23**



To a solution of 4-amino-5-methoxy-3-methylbenzothiophene-2-methanol **22** (180 mg, 0.81 mmol) in acetone (48 mL) was added a solution of potassium nitrosodisulfonate (889 mg, 3.26 mmol) in sodium dihydrogen phosphate buffer (0.3 M; 39 mL). The reaction was stirred at room temperature for 1 h. The excess acetone was removed *in vacuo*. The resulting residue was extracted with dichloromethane and the combined organic layer concentrated. The residue obtained was stirred at room temperature in a 1:1 mixture of hydrochloric acid (2 M)-acetone for 1 h. The acetone was removed *in vacuo*. The resulting residue was extracted with dichloromethane. The organic layer was washed with water, dried over MgSO_4 , filtered and evaporated under reduced pressure to yield the *title compound* (130 mg, 67%) as an orange crystalline solid; mp 215-216 °C; (Found: MH^+ , 239.0379. $\text{C}_{11}\text{H}_{10}\text{O}_4\text{S} + \text{H}$ requires 239.0378); λ_{max} (acetonitrile)/nm 200 (log ϵ 3.86), 232 (3.88), 284 (3.90), 352 (3.24), 412 (3.09); ν_{max} (KBr)/ cm^{-1} 3385, 3054, 2920, 2843, 1677, 1638, 1600, 1538, 1461, 1442, 1342, 1327, 1253, 1227, 1134; δ_{H} (300 MHz; CDCl_3) 5.96 (1 H, s, 6-H), 4.87 (2 H, s, CH_2OH), 3.87 (3 H, s, OMe), 2.47 (3 H, s, Me), 1.71 (1 H, br s, OH); δ_{C} (100 MHz; CDCl_3) 180.0 (C), 176.9


(C), 160.4 (C), 146.6 (C), 143.2 (C), 136.5 (C), 135.7 (C), 107.7 (CH), 58.2 (CH₂), 56.7 (Me), 13.2 (Me); *m/z* (CI) 239 (MH⁺, 100%), 221 (30).

Methyl 3-chloro-5-methoxy-4-nitrobenzothiophene-2-carboxylate **25**



Methyl 3-chloro-5-methoxybenzothiophene-2-carboxylate **24**³¹ (1.00 g, 3.90 mmol) was dissolved in acetic acid (20 mL). Fuming nitric acid (0.25 mL) was added carefully, and the mixture was stirred at room temperature before being heated under reflux overnight. After cooling, the mixture was poured into water (125 mL), and the precipitate collected, washed with water and dried to give the *title compound* (1.03 g, 87%) used without further purification. A sample was recrystallized from methanol to give yellow crystals, mp 170-173 °C; (Found: M⁺, 300.9821. C₁₁H₈³⁵ClNO₅S requires 300.9812); ν_{\max} (KBr)/cm⁻¹ 1721, 1607, 1534, 1514, 1438, 1374, 1293, 1241; δ_{H} (300 MHz; CDCl₃) 7.86 (1 H, d, *J* 9.0, ArH), 7.33 (1 H, d, *J* 9.0, ArH), 3.99 (3 H, s, OMe), 3.97 (3 H, s, OMe); δ_{C} (100 MHz; CDCl₃) 160.8 (C), 149.4 (C), 132.1 (C), 130.1 (C), 127.4 (C), 125.4 (CH), 122.8 (C), 114.6 (CH), 57.4 (Me), 52.9 (Me); 1 C unobserved; *m/z* (EI) 303/301 (MH⁺, 100%).

Methyl 4-amino-5-methoxybenzothiophene-2-carboxylate **26**




The 3-chloro-4-nitrobenzothiophene **25** (1.00 g, 3.3 mmol) and sodium acetate (0.60 g, 7.3 mmol) were dissolved in methanol (200 mL) and THF (55 mL). Palladium-on-carbon (5%; 0.5 g) was added, and the mixture was shaken under a hydrogen atmosphere for 5 h. The mixture was filtered through Celite, the filtrate evaporated, and the residue purified by

chromatography eluting with light petroleum / ethyl acetate (4:1) to give the *title compound* (0.69 g, 88%) as a pale yellow solid, mp 149-151 °C; (Found: C, 55.6; H, 4.7; N, 5.7.


$C_{11}H_{11}NO_3S$ requires C, 55.7; H, 4.7; N, 5.9%); ν_{\max} (KBr)/ cm^{-1} 3445, 3359, 1702, 1625, 1528, 1470, 1206; δ_H (300 MHz; $CDCl_3$) 8.01 (1 H, d, J 0.8, 3-H), 7.20 (1 H, dd, J 8.6, 0.8, ArH), 7.09 (1 H, d, J 8.6, ArH), 4.27 (2 H, br s, NH_2), 3.93 (3 H, s, OMe), 3.91 (3 H, s, OMe); δ_C (100 MHz; $CDCl_3$) 163.3 (C), 142.9 (C), 136.0 (C), 132.4 (C), 131.8 (C), 127.8 (C), 126.5 (CH), 113.7 (CH), 111.4 (CH), 56.8 (Me), 52.3 (Me); m/z (EI) 237 (M^+ , 70%), 222 (100), 194 (35), 83 (68).

Methyl 5-methoxy-4,7-dioxobenzothiophene-2-carboxylate **27**




A mixture of 4-aminobenzothiophene **26** (50 mg, 0.21 mmol) and potassium nitrosodisulfonate (230 mg, 0.84 mmol) in acetone (7.5 mL) and sodium dihydrogen phosphate buffer (0.3 M; 7.5 mL) was stirred at room temperature overnight. The acetone was evaporated, water (70 mL) was added, and the mixture was extracted with dichloromethane (3 × 25 mL). The combined extracts were dried (Na_2SO_4), evaporated and the residue purified by chromatography eluting with dichloromethane to give the *title compound* (40 mg, 75%) as a dark yellow solid, mp 234 °C; (Found: C, 52.8; H, 3.3. $C_{11}H_8O_5S$ requires C, 52.4; H, 3.2%); ν_{\max} (KBr) 3088, 2960, 1720, 1689, 1633, 1597, 1534, 1439, 1244, 1268, 1085; δ_H (300 MHz; $CDCl_3$) 8.15 (1 H, s, 3-H), 6.08 (1 H, s, 6-H), 3.94 (3 H, s, OMe), 3.90 (3 H, s, OMe); δ_C (75 MHz; $CDCl_3$) 179.6 (C), 175.1 (C), 161.4 (C), 160.8 (C), 148.0 (C), 139.3 (C), 138.5 (C), 130.7 (CH), 109.0 (CH), 57.0 (Me), 53.1 (Me); m/z (EI) 252 (M^+ , 100%), 237 (34).

4-Amino-5-methoxybenzothiophene-2-methanol **28**



A solution of the benzothiophene **26** (2.00 g, 8.43 mmol) in dry THF (28 mL) was added dropwise to lithium aluminum hydride (1.07 g, 28.1 mmol) in THF (55 mL). The mixture was stirred at room temperature for 16 h, and then water was carefully added. The mixture was filtered through Celite, and extracted with ethyl acetate (3 × 30 mL). The combined extracts were washed with water (30 mL), dried (MgSO₄), and evaporated to a solid that was recrystallized from methanol to give the *title compound* (1.38 g, 78%), mp 130-132 °C; (Found: C, 57.2; H, 5.3; N, 6.6. C₁₀H₁₁NO₂ requires C, 57.4; H, 5.3; N, 6.7%); ν_{\max} (KBr)/cm⁻¹ 3401, 3310, 3016, 1602, 1480, 1463; δ_{H} (300 MHz; CDCl₃) 7.19 (1 H, d, *J* 8.6, ArH), 7.09 (1 H, s, 3-H), 6.95 (1 H, d, *J* 8.6, ArH), 4.87 (2 H, s, CH₂), 3.90 (3 H, s, OMe); OH, NH₂ not observed; δ_{C} (75 MHz; CDCl₃) 144.0 (C), 143.3 (C), 133.7 (C), 130.4 (C), 128.6 (C), 117.1 (CH), 111.9 (CH), 110.7 (CH), 61.0 (CH₂), 56.8 (Me); *m/z* (EI) 209 (M⁺, 62%), 194 (100), 166 (25).


2-Hydroxymethyl-5-methoxybenzothiophene-4,7-dione **29**



A mixture of the 4-aminobenzothiophene **28** (0.32 g, 1.5 mmol) and potassium nitrosodisulfonate (1.61 g, 6.0 mmol) in acetone (47 mL) and sodium dihydrogen phosphate buffer (0.3 M; 47 mL) was stirred at room temperature for 1 h. Work-up as described above gave the *title compound* (0.29 g, 88%) as a dark orange solid, mp 213 °C (from methanol); (Found: C, 53.3; H, 3.3. C₁₀H₈O₄S requires C, 53.6; H, 3.60%); ν_{\max} (KBr)/cm⁻¹ 3413, 3062, 2986, 2947, 1680, 1628, 1598, 1572, 1326, 1246, 1140, 1084, 1043, 864, 792; δ_{H} (300 MHz; CDCl₃) 7.43 (1 H, s, 3-H), 5.99 (1 H, s, 6-H), 4.91 (2 H, d, *J* 6.0, CH₂OH), 3.88 (3 H, s,

OMe), 2.05 (1 H, t, J 6.0, CH₂OH); δ_C (100 MHz; CDCl₃) 180.2 (C), 175.9 (C), 160.6 (C), 156.3 (C), 142.3 (C), 139.2 (C), 121.5 (CH), 108.7 (CH), 58.8 (CH₂), 57.3 (Me); m/z (EI) 224 (M⁺, 100%) 209 (28), 194 (35), 125 (26).


4-Amino-5-methoxybenzothiophene-2-carboxaldehyde **30**



A solution of the benzothiophene-2-methanol **28** (0.90 g, 4.30 mmol) in dichloromethane (100 mL) was stirred with manganese(IV) oxide (3.78 g, 43 mmol) under reflux for 24 h. The solution was filtered through Celite, washed through with dichloromethane, and the combined filtrate and washings concentrated under vacuum. The residue was purified by chromatography eluting with light petroleum/ethyl acetate (4:1) to give the *title compound* (0.29 g, 33%) as a pale yellow solid, mp 128-130 °C; (Found: C, 57.9; H, 4.3; N, 6.5.

C₁₀H₉NO₂S requires C, 57.9; H, 4.4; N, 6.8%); ν_{\max} (KBr)/cm⁻¹ 3474, 3373, 2826, 1667, 1520, 1483; δ_H (300 MHz; CDCl₃) 10.00 (1 H, s, CHO), 7.97 (1 H, s, H-3), 7.19 (1 H, d, J 8.6, ArH), 7.09 (1 H, d, J 8.6, ArH), 4.42 (2 H, br s, NH₂), 3.90 (3 H, s, OMe); δ_C (75 MHz; CDCl₃) 184.4 (CH), 143.0 (C), 142.2 (C), 136.2 (C), 133.2 (C), 130.8 (CH), 127.7 (C), 114.6 (CH), 111.9 (CH), 56.7 (Me); m/z (EI) 207 (M⁺, 70%), 192 (100), 164 (40), 77 (35).


2-Formyl-5-methoxybenzothiophene-4,7-dione **31**



A mixture of the 4-aminobenzothiophene **30** (50 mg, 0.24 mmol) and potassium nitrosodisulfonate (260 mg, 0.96 mmol) in acetone (7.5 mL) and sodium dihydrogen phosphate buffer (0.3 M; 7.5 mL) was stirred at room temperature overnight. Work-up as


described above gave the *title compound* (42 mg, 79%) as a dark orange solid, mp 218 °C; (Found: C, 53.9; H, 2.8. C₁₀H₆O₄S requires C, 54.0; H, 2.7%); ν_{\max} (KBr)/cm⁻¹ 3070, 1688, 1649, 1600, 1522, 1325, 1249, 1148, 1079, 867; δ_{H} (300 MHz; CDCl₃) 10.03 (1 H, s, CHO), 8.16 (1 H, s, 3-H), 6.12 (1 H, s, 6-H), 3.92 (3 H, s, OMe); δ_{C} (75 MHz; CDCl₃) 183.0 (CH), 179.4 (C), 174.9 (C), 161.0 (C), 149.6 (C), 147.6 (C), 138.7 (C), 132.6 (CH) 109.2 (CH), 57.1 (Me); m/z (EI) 222 (M⁺, 100%), 207 (32), 192 (37).

2-Acetoxymethyl-5-methoxybenzothiophene-4,7-dione **32**



Acetic anhydride (2.7 mL) was added to a solution of the alcohol **29** (75 mg, 0.33 mmol) in pyridine (16 mL), and the mixture was stirred overnight at room temperature. The mixture was diluted with water (27 mL), and extracted with dichloromethane (3 × 50 mL). The combined extracts were washed with water (50 mL) and brine (50 mL), dried (MgSO₄) and evaporated. The residue was purified by chromatography eluting with dichloromethane/ethyl acetate (19:1) to give the *title compound* as a yellow solid (58 mg, 66%), mp 153-155 °C; (Found: C, 54.0; H, 3.7. C₁₂H₁₀O₅S requires C, 54.1; H, 3.8%); ν_{\max} (KBr)/cm⁻¹ 3091, 2983, 2963, 1725, 1677, 1642, 1602, 1534, 1467, 1438, 1333, 1232, 1141, 1085, 1030, 962, 854, 798; δ_{H} (300 MHz; CDCl₃) 7.49 (1 H, s, 3-H), 5.99 (1 H, s, 6-H), 5.26 (2 H, s, CH₂), 3.88 (3 H, s, OMe), 2.13 (3 H, s, Me); δ_{C} (75 MHz; CDCl₃) 179.7 (C), 175.6 (C), 170.3 (C), 160.3 (C), 146.0 (C), 144.6 (C), 138.7 (C), 125.5 (CH), 108.5 (CH), 60.2 (CH₂), 56.8 (Me), 20.7 (Me); m/z (EI) 266 (M⁺, 32%), 251 (8), 224 (100), 207 (36), 178 (20).

5-Methoxy-2-methylbenzothiophene-4,7-dione **33**




(a) A stirred solution of the alcohol **29** (200 mg, 0.89 mmol) in dichloromethane (11 mL) at $-10\text{ }^{\circ}\text{C}$ was treated with triethylamine (140 mg, 0.19 mL, 1.34 mmol). After 10 min, methanesulfonyl chloride (110 mg, 0.08 mL, 0.98 mmol) was added, and the mixture allowed to stir at room temperature overnight. Water (11 mL) and dichloromethane (30 mL) were added, the organic layer separated, washed with water (50 mL), brine (50 mL), dried (MgSO_4) and evaporated to give the *mesylate* as a yellow solid, used without any purification.

(b) The above product was dissolved in THF (11 mL) and added to lithium aluminum hydride (330 mg, 8.63 mmol) in THF (9 mL). The mixture was stirred overnight at room temperature, before the careful addition of water. The mixture was filtered through Celite, and the filtrate extracted with ethyl acetate ($3 \times 50\text{ mL}$). The combined extracts were washed with saturated aqueous sodium hydrogen carbonate (10 mL), water (10 mL), dried (MgSO_4) and evaporated. The residue was purified by chromatography to give the *title compound* (9 mg, 5%) as a yellow solid, mp $217\text{--}220\text{ }^{\circ}\text{C}$; (Found: M^+ , 208.0194. $\text{C}_{10}\text{H}_8\text{O}_3\text{S}$ requires 208.0194); ν_{max} (KBr)/ cm^{-1} 3066, 2923, 1680, 1640, 1596, 1535, 1468, 1329, 1246, 1083, 968, 858, 792; δ_{H} (300 MHz; CDCl_3) 7.22 (1 H, s, 3-H), 5.94 (1 H, s, 6-H), 3.86 (3 H, s, OMe), 2.56 (3 H, s, Me); δ_{C} (75 MHz; CDCl_3) 179.9 (C), 175.9 (C), 160.0 (C), 148.5 (C), 142.4 (C), 139.4 (C), 124.1 (CH), 108.3 (CH), 56.7 (Me), 15.9 (Me); m/z (EI) 208 (M^+ , 64%), 193 (21), 178 (30), 149 (47), 71 (44), 69 (100).


3.4 Synthesis of Indazolequinones

5,7-Dimethoxy-3-methylindazole **35**



To a solution of 3,5-dimethoxyacetophenone **34** (1.58 g, 8.81 mmol) in dry dichloromethane (44 mL) were added bis(trichloroethyl) azodicarboxylate (3.35 g, 8.81 mmol) and $\text{BF}_3 \cdot \text{EtO}$ (539 μl , 4.40 mmol). The mixture was stirred overnight at room temperature, quenched with aqueous ammonium acetate solution (25%; 70 mL) and extracted with ethyl acetate (4×70 mL). The organic layer was dried over Na_2SO_4 , filtered and evaporated under reduced pressure. The crude product obtained was purified by flash chromatography. A mixture of hydrazine intermediate and starting material (20:1) was obtained. To a solution of the above product in glacial acetic acid (49 mL) was added zinc dust (4.94 g, 75.55 mmol). The mixture was stirred at room temperature for 1 h and water (50 mL) followed by aqueous sodium hydroxide (1 M) were added until $\text{pH} = 10$. The mixture was extracted with ethyl acetate. The organic layer was dried over Na_2SO_4 , filtered and evaporated under reduced pressure. The crude product was purified by flash chromatography, eluting with ethyl acetate-light petroleum 1:1 to yield the *title compound* (590 mg; 35%) as a beige solid; mp 154-156 °C (lit.,³² mp 155-156 °C); δ_{H} (300 MHz; CDCl_3) 11.72 (1 H, br s, NH), 6.64 (1 H, d, J 2.3, ArH), 6.44 (1 H, t, J 2.3, ArH), 3.95 (3 H, s, OMe), 3.80 (3 H, s, OMe), 2.45 (3 H, s, Me).


5,7-Dimethoxy-3-methyl-4-nitroindazole **36**



To a solution of 5,7-dimethoxy-3-methylindazole **35** (200 mg, 1.04 mmol) in acetic acid (8 mL), cooled to 0-5 °C, was added a mixture of nitric acid (69 μl , 1.04 mmol) in acetic acid (1 mL). The mixture was stirred at room temperature for 1 h. The reaction was quenched by addition of brine, extracted with ethyl acetate, dried over MgSO_4 , filtered, evaporated and azeotroped with toluene to remove the acetic acid. The crude product obtained was purified by flash chromatography, eluting with ethyl acetate/light petroleum (1:1), to yield the *title*

compound (201 mg, 81%) as a bright yellow crystalline solid, recrystallized from dichloromethane-pentane; mp 219-220 °C; (Found: C, 50.7; H, 4.5; N, 17.6. C₁₀H₁₁N₃O₄ requires C, 50.6; H, 4.7; N, 17.7%); (Found: MH⁺, 238.0829. C₁₀H₁₁N₃O₄ + H requires 238.0828); ν_{\max} (KBr)/cm⁻¹ 3398, 3149, 2917, 1597, 1517, 1308, 1223, 1115, 1059; δ_{H} (300 MHz; CDCl₃) 13.43 (1 H, s, NH), 6.53 (1 H, s, ArH), 4.05 (3 H, s, OMe), 4.00 (3 H, s, OMe), 2.50 (3 H, s, Me); δ_{C} (100 MHz; d₆-DMSO) 148.9 (C), 148.2 (C), 138.6 (C), 129.0 (C), 124.3 (C), 114.7 (C), 94.8 (CH), 58.3 (Me), 56.6 (Me), 13.2 (Me); *m/z* (CI) 238 (MH⁺, 95%), 221 (55), 207 (100), 192 (80).


5,7-Dimethoxy-1,3-dimethyl-4-nitroindazole 37



To a stirred solution of 5,7-dimethoxy-3-methyl-4-nitroindazole **36** (116 mg, 0.60 mmol) in DMSO (2 mL) was added potassium hydroxide (135 mg, 2.42 mmol). The mixture was stirred at room temperature for 30 min and iodomethane (150 μ l, 2.42 mmol) was added dropwise to the solution. The reaction mixture was then stirred at room temperature for a further 4 h. The crude mixture was diluted with ethyl acetate, washed thoroughly with hydrochloric acid (2 M), dried over MgSO₄, filtered and evaporated. The crude product obtained was purified by flash chromatography, eluting with ethyl acetate/light petroleum (1:1), to yield the *title compound* (71 mg, 49%) as a bright yellow crystalline solid, recrystallized from dichloromethane-pentane; mp 168-169 °C; (Found: C, 52.8; H, 5.1; N, 16.9. C₁₁H₁₃N₃O₄ requires C, 52.6; H, 5.2; N, 16.7%); (Found: M⁺, 251.0910. C₁₁H₁₃N₃O₄ requires 251.0906); ν_{\max} (KBr)/cm⁻¹ 3435, 2929, 2841, 1577, 1522, 1324, 1280, 1220, 1050; δ_{H} (300 MHz; CDCl₃) 6.45 (1 H, s, ArH), 4.19 (3 H, s, NMe), 4.02 (3 H, s, OMe), 3.97 (3 H, s, OMe), 2.40 (3 H, s, Me); δ_{C} (75 MHz; CDCl₃) 149.0 (C), 148.6 (C), 138.9 (C), 128.9 (C),


126.3 (C), 117.1 (C), 94.6 (CH), 58.9 (Me), 56.4 (Me), 39.0 (Me), 13.4 (Me); m/z (EI) 251 (M^+ , 100%), 234 (28), 221 (29), 206 (72), 175 (40). No nOe enhancement observed after pre-irradiation of the Me group at 2.39 ppm. 4.9% nOe enhancement observed on the aromatic proton at 6.45 ppm after pre-irradiation at 4.02 ppm (5-OMe). 2.3% nOe enhancement observed at 3.97 ppm (7-OMe) after pre-irradiation of NMe at 4.19 ppm. 3.5% nOe enhancements observed on the aromatic proton at 6.45 ppm and 1.3% nOe enhancement observed at 4.02 ppm (5-OMe) after pre-irradiation at 3.97 ppm (7-OMe).

4-Amino-5,7-dimethoxy-1,3-dimethylindazole 38




To a suspension of 5,7-dimethoxy-1,3-dimethyl-4-nitroindazole **37** (55.3 mg, 0.22 mmol) in ethanol (3.7 mL) were added tin powder (120 mg, 0.99 mmol) and hydrochloric acid (3 M; 1.5 mL). The mixture was heated under reflux for 1 h. Upon cooling, the solution was decanted from the excess tin and neutralized with a saturated aqueous solution of sodium hydrogen carbonate. The precipitate obtained was extracted with ethyl acetate. The organic layer was filtered through a pad of Celite, dried over $MgSO_4$, filtered and evaporated under reduced pressure to yield the *title compound* (47 mg; 96%) as a colorless solid; mp 79-80 °C; (Found: M^+ , 221.1163. $C_{11}H_{15}N_3O_2$ requires 221.1164); ν_{max} (KBr)/ cm^{-1} 3434, 3343, 2927, 2836, 1590, 1527, 1453, 1351, 1271, 1197, 1111; δ_H (300 MHz; $CDCl_3$) 6.48 (1 H, s, 6-H), 4.11 (3 H, s, NMe), 3.84 (3 H, s, OMe), 3.82 (3 H, s, OMe), 2.66 (3 H, s, Me); δ_C (75 MHz; $CDCl_3$) 139.9 (C), 138.4 (C), 138.3 (C), 130.5 (C), 123.2 (C), 116.3 (C), 99.3 (CH), 59.2 (Me), 56.4 (Me), 38.3 (Me), 14.5 (Me); m/z (EI) 221 (M^+ , 27%), 206 (100), 191 (14), 163 (8).

5-Methoxy-1,3-dimethylindazole-4,7-dione 39



To a solution of 4-amino-5,7-dimethoxy-1,3-dimethylindazole **38** (0.250 g, 1.15 mmol) in acetone (70 mL) was added a solution of potassium nitrosodisulfonate (1.250 g, 4.59 mmol) in sodium dihydrogen phosphate buffer (0.3 M, 58 mL). The reaction was stirred at room temperature for 1h. The excess acetone was removed in *vacuo*. The resulting residue was extracted with dichloromethane. The organic layer was washed with water, dried over MgSO₄, filtered, evaporated under reduced pressure to yield the *title compound* (208 mg, 78%) as a bright yellow crystalline solid, recrystallized from dichloromethane-pentane; mp 185-186 °C; (Found: C, 58.2; H, 4.8; N, 13.6. C₁₀H₁₀N₂O₃ requires C, 58.2; H, 4.9; N, 13.6%); (Found: M⁺, 206.0683. C₁₀H₁₀N₂O₃ requires 206.0691); λ_{max} (acetonitrile)/nm 272 (log ε 3.91), 320 (3.84); ν_{max} (KBr)/cm⁻¹ 1680, 1657, 1590, 1529, 1510, 1340, 1216, 1018; δ_H (300 MHz; CDCl₃) 5.79 (1 H, s, 6-H), 4.16 (3 H, s, NMe), 3.85 (3 H, s, OMe), 2.49 (3 H, s, Me); δ_C (75 MHz; CDCl₃) 177.8 (C), 176.1 (C), 162.0 (C), 149.0 (C), 137.4 (C), 117.1 (C), 107.3 (CH), 57.0 (Me), 38.4 (Me), 12.9 (Me); *m/z* (EI) 206 (M⁺, 100%), 191 (85), 177 (62), 123 (30).

3-Hydroxymethyl-5-methoxy-1-methylindazole-4,7-dione **40**




A solution of 5-methoxy-1,3-dimethylindazole-4,7-dione **39** (51 mg, 0.25 mmol), AIBN (12 mg, 0.07 mmol) and *N*-bromosuccinimide (86 mg, 0.49 mmol) in CCl₄ (3 mL) was purged five times with vacuum followed with nitrogen. The reaction mixture was heated at reflux under a nitrogen atmosphere overnight. The mixture was concentrated under reduced pressure

and purified by flash chromatography, eluting with dichloromethane, to yield the *bromide intermediate*, which was used in the next step with no further purification. The *bromide intermediate* dissolved in acetone (9 mL) and water (4 mL) was added to a suspension of silver nitrate (92 mg, 0.54 mmol) in an aqueous solution of acetone (50%; 13 mL). The reaction mixture was heated under reflux overnight and extracted with ethyl acetate. The combined organic layer was dried over MgSO₄, filtered and evaporated under reduced pressure. The residue obtained was purified by flash chromatography, eluting with light petroleum-ethyl acetate 1:2, to yield the *title compound* (29 mg; 36%) as a yellow solid; mp 177-179 °C; (Found: MH⁺, 223.0721. C₁₀H₁₀N₂O₄ + H requires 223.0719); λ_{max} (acetonitrile)/nm 276 (log ε 4.23), 308 (3.74); ν_{max} (KBr)/cm⁻¹ 3368, 2928, 1689, 1653, 1593, 1525, 1505, 1240, 1204, 1015; δ_H (300 MHz; CDCl₃) 5.87 (1 H, s, 6-H), 4.86 (2 H, s, CH₂OH), 4.17 (3 H, s, NMe), 3.90 (3 H, s, OMe); δ_C (75 MHz; CDCl₃) 177.5 (C), 177.0 (C), 161.8 (C), 152.6 (C), 137.8 (C), 117.4 (C), 107.7 (CH), 58.2 (CH₂), 57.2 (Me), 38.7 (Me); *m/z* (ES) 223 (MH⁺, 95%), 164 (35), 149 (30), 90 (100).

3.5 Synthesis of Benzisoxazolequinones


(*E*)-2-Hydroxy-5-methoxyacetophenone oxime **42**



To a solution of 2-hydroxy-5-methoxyacetophenone **41** (2.50 g, 15.1 mmol) and hydroxylamine hydrochloride (2.10 g, 30.1 mmol) in ethanol (50 mL) was added dropwise dry pyridine (2.68 mL, 30.1 mmol). The reaction mixture was stirred at room temperature overnight and evaporated under reduced pressure. The crude product obtained was purified by flash chromatography, eluting with ethyl acetate-light petroleum 1:2, to yield the *title compound* (1.68 g, 98%) as a colorless solid; mp 112-113 °C (lit.,³⁵ mp 121 °C); (Found: MH⁺, 182.0813. C₉H₁₁NO₃ + H requires 182.0817); ν_{max} (KBr)/cm⁻¹ 3348, 2964, 2920, 2831,


1641, 1497, 1404, 1368, 1284, 1232, 1208, 1176, 1052, 1011; δ_{H} (300 MHz; CDCl_3) 10.70 (1 H, br s, OH), 7.53 (1 H, br s, OH), 7.17 (1 H, d, J 3.0, 6-H), 7.11 (1 H, dd, J 9.0, 3.0, 4-H), 6.93 (1 H, d, J 9.0, 3-H), 3.80 (3 H, s, OMe), 2.62 (3 H, s, Me); δ_{C} (100 MHz; CDCl_3) 159.3 (C), 152.3 (C), 151.6 (C), 118.7 (C), 117.7 (CH), 116.5 (CH), 113.1 (CH), 56.0 (Me), 10.9 (Me); m/z (CI) 182 (MH^+ , 63%), 166 (100), 151 (15), 125 (5).

5-Methoxy-3-methylbenzisoxazole 43




Diisopropyl azodicarboxylate (8.5 mL, 43.5 mmol) was added dropwise to a stirred solution of the oxime **42** (5.24 g, 29.0 mmol) and triphenylphosphine (11.35 g, 43.5 mmol) in dry THF (192 mL) under nitrogen. The reaction mixture was stirred at room temperature overnight, and evaporated under reduced pressure. The crude product was purified by chromatography, eluting with ethyl acetate/light petroleum (1:3), to yield the *title compound* (4.72 g, 100%) as a yellow solid; mp 28-30 °C (lit.,³⁶ mp not given); (Found: MH^+ , 164.0708. $\text{C}_9\text{H}_9\text{NO}_2 + \text{H}$ requires 164.0712); ν_{max} (KBr)/ cm^{-1} 2972, 2935, 1730, 1528, 1484, 1458, 1439, 1252, 1219, 1127, 1079, 1028; δ_{H} (300 MHz; CDCl_3) 7.44 (1 H, d, J 9.0, 7-H), 7.16 (1 H, dd, J 9.0, 2.3, 6-H), 6.96 (1 H, d, J 2.3, 4-H), 3.87 (3 H, s, OMe), 2.56 (3 H, s, Me); δ_{C} (75 MHz; CDCl_3) 158.4 (C), 156.2 (C), 154.9 (C), 122.5 (C), 120.2 (CH), 110.5 (CH), 101.3 (CH), 55.9 (Me), 10.1 (Me); m/z (CI) 164 (MH^+ , 100%), 148 (5).

5-Methoxy-3-methyl-4-nitrobenzisoxazole 44



To a solution of nitric acid/sulfuric acid 9:1 (22 mL), cooled in a salt and ice bath, was added 5-methoxy-3-methylbenzoxazole **43** (2.22 g, 13.6 mmol) portionwise. The mixture was stirred at room temperature overnight. The mixture was poured into ice/water, basified with a saturated aqueous sodium hydrogen carbonate, extracted with dichloromethane, dried over MgSO₄, and evaporated under reduced pressure to yield the *title compound* (2.58 g, 91%) as a yellow solid; mp 103-105 °C; (Found: MH⁺, 209.0551. C₉H₈N₂O₄ + H requires 209.0562); ν_{\max} (KBr)/cm⁻¹ 3092, 2954, 2850, 1525, 1513, 1475, 1371, 1325, 1267, 1060; δ_{H} (300 MHz; CDCl₃) 7.69 (1 H, d, *J* 9.2, ArH), 7.36 (1 H, d, *J* 9.2, ArH), 4.00 (3 H, s, OMe), 2.50 (3 H, s, Me); δ_{C} (75 MHz; CDCl₃) 158.2 (C), 153.0 (C), 149.1 (C), 131.7 (C), 116.9 (CH), 115.3 (C), 113.6 (CH), 58.1 (Me), 10.9 (Me); *m/z* (CI) 209 (MH⁺, 100%), 192 (5).

5-Methoxy-3-methylbenzoxazole-4,7-dione **45**



To a suspension of 5-methoxy-3-methyl-4-nitrobenzoxazole **44** (2.5 g, 12.0 mmol) in ethanol (200 mL) was added tin powder (3.6 g, 120.2 mmol) and hydrochloric acid (3 M; 83 mL). The mixture was stirred and heated under reflux for 1 h. Upon cooling, the reaction mixture was decanted from the excess of tin and neutralized with a saturated sodium hydrogen carbonate. The suspension obtained was added to an equal volume of water. The precipitate and aqueous layer were stirred overnight with dichloromethane, filtered through Celite and the layers separated. The organic layer was dried over Na₂SO₄ and concentrated to yield the 4-amino compound, used in the next step with no further purification.

To a solution of the 4-amino compound in acetone (490 mL) was added a solution of potassium nitrosodisulfonate (8.9 g, 33.2 mmol) in sodium dihydrogen phosphate buffer (0.3 M; 400 mL). The reaction was stirred at room temperature for 1 h, and concentrated under

reduced pressure. The residue was extracted with dichloromethane. The organic layer was washed with water, dried over MgSO_4 , filtered, evaporated under reduced pressure to yield the *title compound* (250 mg, 20%) as a dark orange crystalline solid; mp 126-129 °C; (Found: MH^+ , 194.0451. $\text{C}_9\text{H}_7\text{NO}_4 + \text{H}$ requires 194.0453); λ_{max} (acetonitrile)/nm 220 (log ϵ 3.92), 256 (3.89), 380 (3.30); ν_{max} (KBr)/ cm^{-1} 3437, 3060, 2917, 1698, 1667, 1578, 1486, 1467, 1440, 1355, 1336, 1251, 1205, 1186, 1032, 1013; δ_{H} (300 MHz; CDCl_3) 5.98 (1 H, s, 6-H), 3.93 (3 H, s, OMe), 2.59 (3 H, s, Me); δ_{C} (75 MHz; CDCl_3) 176.0 (C), 174.5 (C), 165.2 (C), 161.8 (C), 158.1 (C), 117.7 (C), 107.1 (CH), 57.9 (Me), 11.0 (Me); m/z (CI) 194 (MH^+ , 100%), 166 (5).

3.6 Biology

HPLC analysis

Reduction of the quinones was followed by HPLC using an Alltech C18 (5 μm , 250 mm \times 4.6 mm) column with a Waters HPLC system (2487 Dual λ Absorbance detector, two 515 HPLC pumps, 717plus Autosampler, Millennium32 Chromatography Manager). The solvent program used a linear gradient of 5% to 80% B over 10 min, 80% B for 5 min, then 80% B to 5% B over 5 min (solution A, 10 mM potassium phosphate buffer, pH 6.0; solution B, methanol). Reactions were run in 25 mM Tris-HCl (pH 7.4) containing 200 μM NADH (Sigma), 50 μM quinone, and recombinant human NQO1 (gift from David Ross, University of Colorado, Denver, CO). NADH oxidation was quantified at 340 nm following 30-40 min incubations at 22 °C.

Spectrophotometric method

Quinone reduction by recombinant human NQO1 was also quantified using a modification of an assay that uses cytochrome *c* as the terminal electron acceptor.³⁷ Reaction mixtures

contained 1 mM NADH (Sigma), 25 μ M quinone, 70 μ M cytochrome *c* (Sigma) and 0.1-3.0 μ g/ml rhNQO1 in 25 mM Tris-HCl (pH 7.4) with 0.07% BSA and 0.1% Tween-20. Reactions were run at least in triplicate at 22 °C in a Beckman DU 7500 spectrophotometer at 550 nm (molar absorptivity 21.1 mM⁻¹ cm⁻¹ for cytochrome *c*). Initial reduction rates (μ mol cytochrome *c* reduced/min/mg NQO1) were calculated from the linear portion (0-30 s) of the reaction curves.

Acknowledgements

This work was supported by the FORCE Cancer Charity, the Cancer Research Campaign (as was) [CRUK], and NIH Grant P20RR017670 (H.D.B.). We also thank the EPSRC Mass Spectrometry Centre at Swansea for mass spectra, Dr Stephen Green for help with the electrochemical experiments, and Professor David Ross (University of Colorado, Denver) for generous gifts of enzyme.

References

1. Thomson, R. H. *Naturally Occurring Quinones IV. Recent advances*, 4th ed.; Blackie: London, 1997.
2. Tisler, M. *Adv. Heterocycl. Chem.* **1989**, *45*, 37-150.
3. Boutros, R.; Lobjois, V.; Ducommun, B. *Nat. Rev. Cancer* **2007**, *7*, 495-507.
4. Carter, S. K.; Crooke, S. T. *Mitomycin C; Current Status and New Developments*; Academic Press: New York, 1979.
5. Remers, W. A.; Dorr, R. T. In *Alkaloids: Chemical and Biological Perspectives*; Pelletier, S. W. Ed.; Wiley: New York, 1988; pp. 1-74.
6. Franck, R. W.; Tomasz, M. In *Chemistry of Antitumor Agents*; Wilman, D. E. V. Ed.; Blackie and Son Ltd.: Glasgow, 1990; pp. 379-393.
7. Danishefsky, S. J.; Schkeryantz, J. M. *Synlett* **1995**, 475-490.
8. Bringmann, G.; Reichert, Y.; Kane, V. V. *Tetrahedron* **2004**, *60*, 3539-3574.
9. Ernster, L. *Chem. Scripta* **1987**, *27A*, 1-13.
10. Ross, D.; Siegel, D.; Beall, H.; Prakash, A. S.; Mulcahy, R. T.; Gibson, N. W. *Cancer Metast. Rev.* **1993**, *12*, 83-101.
11. Beall, H. D.; Winski, S. L. *Frontiers in Bioscience* **2000**, *5*, D639-D648.
12. Colucci, M. A.; Couch, G. D.; Moody, C. J. *Org. Biomol. Chem.* **2008**, *6*, 637-656.
13. Beall, H. D.; Winski, S.; Swann, E.; Hudnott, A. R.; Cotterill, A. S.; O'Sullivan, N.; Green, S. J.; Bien, R.; Siegel, D.; Ross, D.; Moody, C. J. *J. Med. Chem.* **1998**, *41*, 4755-4766.
14. Newsome, J. J.; Swann, E.; Hassani, M.; Bray, K. C.; Slawin, A. M. Z.; Beall, H. D.; Moody, C. J. *Org. Biomol. Chem.* **2007**, *5*, 1629-1640.
15. Faig, M.; Bianchet, M. A.; Winski, S.; Hargreaves, R.; Moody, C. J.; Hudnott, A. R.; Ross, D.; Amzel, L. M. *Structure* **2001**, *9*, 659-667.
16. Winski, S. L.; Swann, E.; Hargreaves, R. H. J.; Dehn, D. L.; Butler, J.; Moody, C. J.; Ross, D. *Biochem. Pharmacol.* **2001**, *61*, 1509-1516.
17. Winski, S. L.; Faig, M.; Bianchet, M. A.; Siegel, D.; Swann, E.; Fung, K.; Duncan, M. W.; Moody, C. J.; Amzel, M.; Ross, D. *Biochemistry* **2001**, *40*, 15135-15142.
18. Fryatt, T.; Pettersson, H. I.; Gardipee, W. T.; Bray, K. C.; Green, S. J.; Slawin, A. M. Z.; Beall, H. D.; Moody, C. J. *Bioorg. Med. Chem.* **2004**, *12*, 1667-1687.
19. Newsome, J. J.; Colucci, M. A.; Hassani, M.; Beall, H. D.; Moody, C. J. *Org. Biomol. Chem.* **2007**, *5*, 3665-3673.
20. Skibo, E. B.; Islam, I.; Schulz, W. G.; Zhou, R.; Bess, L.; Boruah, R. *Synlett* **1996**, 297-309.
21. Esser, L.; Quinn, B.; Li, Y. F.; Zhang, M. Q.; Elberry, M.; Yu, L.; Yu, C. A.; Xia, D. *J. Mol. Biol.* **2004**, *341*, 281-302.
22. Scannell, R. T.; Stevenson, R. *J. Org. Chem.* **1983**, *48*, 127-129.
23. De Rosa, M.; De Rosa, S.; Gambacorta, A.; Minale, L.; Thomson, R. H.; Worthington, R. D. *J. Chem. Soc., Perkin Trans. 1* **1977**, 653-657.
24. Li, Z. R.; Shokes, J. E.; Kounosu, A.; Imai, T.; Iwasaki, T.; Scott, R. A. *Biochemistry* **2003**, *42*, 15003-15008.
25. Sucrow, W.; Brockmann, R.; Haupt, H. J.; Preut, H. *Liebigs Ann. Chem.* **1984**, 1711-1718.
26. Berhe, S.; Slupe, A.; Luster, C.; Charlier, H. A., Jr.; Warner, D. L.; Zalkow, L. H.; Burgess, E. M.; Enwerem, N. M.; Bakare, O. *Bioorg. Med. Chem.* **2010**, *18*, 134-141.
27. Struebe, F.; Rath, S.; Mattay, J. *Eur. J. Org. Chem.* **2011**, 4645-4653.
28. Honey, M. A.; Blake, A. J.; Campbell, I. B.; Judkins, B. D.; Moody, C. J. *Tetrahedron* **2009**, *65*, 8995-9001.

29. Frigoli, M.; Mehl, G. H. *Chem. Eur. J.* **2004**, *10*, 5243-5250.
30. Deng, H.; Fang, Y. *ACS Med. Chem. Lett.* **2012**, *3*, 550-554.
31. Anderson, D. R.; Meyers, M. J.; Kurumbail, R. G.; Caspers, N.; Poda, G. I.; Long, S. A.; Pierce, B. S.; Mahoney, M. W.; Mourey, R. J. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 4878-4881.
32. Mitchell, H.; Leblanc, Y. *J. Org. Chem.* **1994**, *59*, 682-687.
33. Swann, E.; Barraja, P.; Oberlander, A. M.; Gardipee, W. T.; Hudnott, A. R.; Beall, H. D.; Moody, C. J. *J. Med. Chem.* **2001**, *44*, 3311-3319.
34. Bernatek, E. *Acta Chem. Scand.* **1953**, *7*, 677-681.
35. Lindemann, H.; Pickert, W. *Liebigs Ann. Chem.* **1927**, *456*, 275-311.
36. Udd, S.; Jokela, R.; Franzen, R.; Tois, J. *Tetrahedron Lett.* **2010**, *51*, 1030-1033.
37. Phillips, R. M.; Naylor, M. A.; Jaffar, M.; Doughty, S. W.; Everett, S. A.; Breen, A. G.; Choudry, G. A.; Stratford, I. J. *J. Med. Chem.* **1999**, *42*, 4071-4080.