LASER and Light Treatments for HS

Lt Col Rachel Howes

Burns, Plastic and Reconstructive Surgeon

Aims

Overview of LASER and Light Therapies

Evidence for the BEST Laser

Targets - choosing different modalities or combining modalities

Strategies to incorporate LASER and light therapy into HS Management

Questions/Discussion


Overview of LASER and Light Therapies

LASER

- Monochromatic, coherent
- Continuous/pulsed/Q-switched

Intense Pulsed light (IPL)

- Xenon flash lamp + filter
- Divergent







Chromophores

- Melanin
 - hair, skin pigmentation
- Water
 - ablates cells
- Oxyhaemoglobin
 - vessels, redness


Laser Absorption



Laser Wavelength (Colour of light)




Depth of Penetration by Wavelength



Evidence for the BEST Laser/Light Treatment

- Laser hair removal using various modalities is safe and effective in the treatment of HS
- Most evidence for Nd:YAG
 - Better for darker skin types
 - Reduces inflammation too
- IPL/Alexandrite/Diode
- +/- CO2
- Variability explained by differing protocols, follow up timelines and outcome measures



Targets in HS


Hair follicle

Inflammation


Abscess


Sinus Tract

Scarring



LASER HAIR REMOVAL





Cycles of treatment (hair growth cycle)
Depth of hair (dermal thickness)
Hair colour and thickness


Aims to stop disease process Secondary benefits of reduced inflammation – not sustained

Inflammation

IPL

- Selective photothermolysis
- Targets both haemoglobin and melanin in and around the hair follicle
- Result = anti-inflammation and follicular destruction

- 420-1,200 nm, 20–100 ms pulse duration, 7– 32 J/cm2.
- Settings vary by skin type and device
- Not recommended in Fitzpatrick skin types IV-VI



CO2 Laser Surgery

Excisional/deroofing procedure (LA)


Continuous mode

Heals by secondary intention (3-5 weeks)

Reduce flares

Address sinus tracts/nodules

Reduces recurrence at site of surgery, not regionally



Sinus tracts


- Diode Laser 1470nm
- PDT intralesional photodynamic therapy with 5-aminolevulinic acid and 630 nm laser
- Inside sinus tracts
- Minimally invasive obliteration of epithelium in the tracts
- Avoids damage to intact overlying skin
- LA procedure
- Treating sinus tract, not underlying disease process (hair follicle)



Scarring

- Fractional CO2 Laser
- Improves surgical scarring if delivered within
 1 month of surgery
- Improves burn-related scarring thickness, pliability
- Improves acne type scarring resurfacing







Lewis CJ, Douglas H, Martin L, Deng Z, Melton P, Fear MW, Wood FM, Rea S. Carbon dioxide laser treatment of burn-related scarring: Results of the ELIPSE (Early Laser Intervention Promotes Scar Evolution) prospective randomized controlled trial. J Plast Reconstr Aesthet Surg. 2023 Sep;84:368-376. doi: 10.1016/j.bjps.2023.06.012. Epub 2023 Jun 19. PMID: 37393760.



Questions/Comments

Thanks very much for your attention