

Dosing table for IV Iron within the POP-I trial\*

Values of Iron (mg) rounded down to the nearest 100mg

	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)
Hb 100-110 (g/L)	500	500	500	500	500	500	500	500	500	500	500	500	500	500	500
Hb<100 (g/L)	500	500	500	500	500	500	500	500	500	500	500	500	500	500	500
Weight (kg)	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39

	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)
Hb 100-110 (g/L)	500	500	500	500	500	500	500	500	500	500	1000	1000	1000	1000	1000
Hb<100 (g/L)	500	500	500	500	500	500	500	500	500	500	1000	1000	1000	1000	1000
Weight (kg)	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54

	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)
Hb 100-110 (g/L)	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Hb<100 (g/L)	1100	1100	1100	1100	1100	1200	1200	1200	1200	1200	1300	1300	1300	1300	1300
Weight (kg)	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69

	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)
Hb 100-110 (g/L)	1400	1400	1400	1400	1400	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500
Hb<100 (g/L)	1400	1400	1400	1400	1400	1500	1500	1500	1500	1500	1600	1600	1600	1600	1600
Weight (kg)	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84

	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)	Dose (mg)
Hb 100-110 (g/L)	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500
Hb<100 (g/L)	1700	1700	1700	1700	1700	1800	1800	1800	1800	1800	1900	1900	1900	1900	1900
Weight (kg)	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99

For patient's whose weight is  $\geq 100$ kg, administer 1500mg (Hb 100g-110g/L) or 2000mg (Hb<100g/L)