

DR LISA KIRBY (Orcid ID : 0000-0002-9231-1927)

DR SONIA GRAN (Orcid ID : 0000-0002-2443-5100)

DR OLUWAFIKUNAYO OREKOYA (Orcid ID : 0000-0002-5945-1995)

DR ROSALIND SIMPSON (Orcid ID : 0000-0002-8093-7965)

Article type : Research Letter

Is urinary incontinence associated with vulval lichen sclerosus in females? A cross-sectional study

L. Kirby,¹ S. Gran,¹ F. Orekoya,² C. Owen,² and R. Simpson¹.

¹Centre of Evidence Based Dermatology, School of Medicine, University of Nottingham, Nottingham; ²Department of Dermatology, Blackburn Hospital, East Lancashire Hospitals NHS Trust, Blackburn.

Correspondence: Lisa Kirby

E-mail: lisa.kirby@nottingham.ac.uk

<https://orcid.org/0000-0002-9231-1927> (L Kirby)

<https://orcid.org/0000-0002-2443-5100> (S Gran)

<https://orcid.org/0000-0002-8093-7965> (R Simpson)

Funding sources: British Skin Foundation (010_BSFBAD_19)

Conflicts of interest: none to declare

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/bjd.20583](https://doi.org/10.1111/bjd.20583)

This article is protected by copyright. All rights reserved

The data that support the findings of this study and a full study report are openly available in University of Nottingham Research Data Management Repository at <http://doi.org/10.17639/nott.7114>

Dear Editor,

Increasing evidence suggests the pathogenesis of male genital lichen sclerosus (LS) is driven by chronic, occluded exposure of susceptible epithelium to the irritant effect of urine (1, 2). It is not clear how this can be extrapolated to females, however, the typical 'figure of eight' distribution seen in female genital LS suggests that urine could be implicated. This has not yet been robustly investigated (3).

This study utilised data prospectively entered into the East Lancashire Hospitals NHS Trust vulval dermatology database between 2017-2020. Cross-sectional analysis was performed to determine the odds of urinary incontinence (UI) in LS, compared with other genital conditions in adult females. The LS group were women with genital LS, diagnosed clinically and/or histologically. The non-LS group were women with genital conditions other than LS, identified from the same clinic. UI was screened for using the validated International Consultation on Incontinence Questionnaire Urinary Incontinence (ICIQ) Short Form (4). Power calculation demonstrated for 90% power and 5% error to detect a 20% difference, 110 patients were needed per group. Ethical approval was not required as data were unlinked and anonymous.

All patients were assessed by a Consultant Dermatologist with a specialist interest in vulval disease. Clinical diagnosis (plus histological if available), demographics, body mass index (BMI), parity and ICIQ score were recorded.

Age was identified as an a priori confounder; BMI and parity were potential confounders(5). The association between potential confounders and LS and UI status were analysed using the two-sample t-test for normally distributed data or Mann-Whitney U-test for non-normally distributed data. Univariate logistic regression was conducted to determine the association between LS and UI. Multivariable logistic

regression using the 10% change in the adjusted odds ratio (OR) was used to assess for confounding and to obtain a final model, which included all confounders. Sensitivity analyses were conducted excluding patients with LS/LP overlap and irritant contact dermatitis (ICD) secondary to UI.

The most common diagnoses in the non-LS group were eczema, candidiasis, vulvodynia and irritant contact dermatitis. Baseline characteristics for all participants (n=384) are presented in table 1. The prevalence of UI was 63% in the LS group and 34% in the non-LS group, $P < 0.001$. Women with LS were significantly older (median 62 versus 50, $P < 0.001$). The LS group had higher BMI (median 29 versus 26 ($P < 0.001$)) and parity (median 2, IQR 2-3 versus 0-2, $P < 0.001$). The LS group reported lower washing frequency than the non-LS group (median 1, IQR 1-1 versus 1-2, $P = 0.047$.)

Parity and BMI were significantly associated with both LS and UI status. Participants with missing data for these variables or UI status were excluded from the final analysis (n=31); 353 participants were included in the multivariable analysis (LS n=112, non-LS n=241). BMI and parity were not found to be confounders using the 10% rule. The unadjusted OR for UI was 3.85 (95%CI 2.40-6.18), $P < 0.001$. The final age-adjusted OR was 2.56 (95%CI 1.55-4.24), $P < 0.001$.

Sensitivity analysis excluding LS/LP overlap (n=8) from the LS group increased the age-adjusted OR to 2.80 (95%CI 1.67-4.70) $P < 0.001$. When ICD secondary to UI (n=21) was excluded from the non-LS group, OR was 3.15 (95%CI 1.87-5.30). Excluding both these groups, the age-adjusted OR was 3.55 (95%CI 2.10- 6.04) $P < 0.001$.

To our knowledge, this is the first study using a validated screening tool for UI and prospectively collected data, which is powered to determine the odds of UI in vulval LS.

The prevalence of UI was 63% and the odds were increased 2.5-fold.

Few studies report the prevalence of UI in women with LS. A recent meta-analysis found pooled prevalence was 0.35 (95%CI 0.13-0.58), comparable with that of the general population(3). Our LS population were similar to those in two previous cross-sectional studies; older with higher parity (5) and BMI than those without LS (6). In women of comparable age and BMI to our LS population, 46% of women aged 60-64 (7) and 50% of those with BMI 25-29 had UI(8). Less frequent washing

in women with LS is an interesting finding, however discussion is beyond the scope of this letter.

The strengths of this study were the use of a validated screening tool for UI, and diagnosis by an expert clinician, with histological confirmation when required. The entire database was utilised to minimise selection bias. Data were collected from a single centre; therefore, study population and secondary care setting may limit generalisability. A key limitation associated with the cross-sectional design is that the direction of association between UI and LS cannot be determined.

This study provides evidence of a link between UI and vulval LS. Large population-based cohort studies are now needed to determine the nature of this association.

Acknowledgments: we would like to thank Mr Pete Sagar for the database design and Mr Simon Hill for advice about evaluating urinary incontinence.

References

1. Bunker CB, Patel N, Shim TN. Urinary voiding symptomatology (micro-incontinence) in male genital lichen sclerosis. *Acta Derm Venereol.* 2013;93(2):246-8.
2. Bunker CB, Shim TN. Male genital lichen sclerosis. *Indian J Dermatol.* 2015;60(2):111-7.
3. Kirby L, Gran, S, Kreuser-Genis, I, Owen, C, Simpson, R. Is urinary incontinence associated with lichen sclerosis in females? A systematic review and meta-analysis. *Skin Health Dis.* 2021.

4. Avery K, Donovan J, Peters TJ, Shaw C, Gotoh M, Abrams P. ICIQ: a brief and robust measure for evaluating the symptoms and impact of urinary incontinence. *Neurourol Urodyn*. 2004;23(4):322-30.

5. Kennedy CM, Nygaard IE, Bradley CS, Galask RP. Bladder and bowel symptoms among women with vulvar disease are they universal? *Journal of Reproductive Medicine for the Obstetrician and Gynecologist*. 2007;52(12):1073-8.
6. Swenson CW, Menees SB, Haefner HK, Berger MB. Lower Urinary Tract and Functional Bowel Symptoms in Women With Vulvar Diseases and Controls. *Female Pelvic Medicine & Reconstructive Surgery*. 2015;21(4):211-4.
7. Hunskaar S, Lose G, Sykes D, Voss S. The prevalence of urinary incontinence in women in four European countries. *BJU Int*. 2004;93(3):324-30.
8. Schreiber Pedersen L, Lose G, Hoybye MT, Elsner S, Waldmann A, Rudnicki M. Prevalence of urinary incontinence among women and analysis of potential risk factors in Germany and Denmark. *Acta Obstet Gynecol Scand*. 2017;96(8):939-48.

Table 1. Baseline characteristics of study participants

Variable	LS group (n=126)	Missing data (LS)	Non-LS group (n=258)	Missing data (non-LS)	Statistical significance (P-value)
Age (median, IQ range)	62 (55-71)	0	50 (34-64)	0	<0.001
BMI (median, IQ range)	29 (25.5-33)	10	26 (23-32)	12	<0.001
Parity (median, IQ range)	2 (2-3)	2	2 (0-2)	2	<0.001
Self-reported UI (number, percentage)	n=79 (63%)	2	n=87 (34%)	3	<0.001
ICIQ score (median, IQ range)	5 (0-11)	4	0 (0-5)	5	<0.001
Washing frequency (median, IQ range)	1(1-1)	0	1(1-2)	6	0.047